CORRECTION Open Access



## Correction to: Transdiagnostic group therapy for people with self-critic and low self esteem, based on compassion focused therapy principles

Bodil Andersen\* and Peter Hørslev Rasmussen

## Correction

Unfortunately, after publication of this article [1], it was noticed that Figs. 2, 3 and 4 were replications of Fig. 1. The correct versions of Figs. 2, 3 and 4 can be seen here.


Received: 25 January 2018 Accepted: 25 January 2018 Published online: 10 February 2018

## Reference

 Andersen B, Rasmussen PH. Transdiagnostic group therapy for people with self-critic and low self esteem, based on compassion focused therapy principles. J Compassionate Health Care. 2017;4:14. https://doi.org/10.1186/ s40639-017-0043-1.


<sup>\*</sup> Correspondence: bodil.andersen@dadlnet.dk Kognitiv Center Fyn, Odense, Denmark






The box designates the median and the upper and lower quartiles. Whiskers/lines represent the max. and min. observations within the range of the 1. and 3. quartiles  $\pm 1.5 \times 10^{10}$  (InterQuartile Range). Dots are outlier cases. Bar plots show change from pre to post measurements for the response variable.

Fig. 2 Change in BDI according to diagnosis



Whiskers/lines represent the max. and min. observations within the range of the 1. and 3. quartiles +/- 1.5\*IQR (InterQuartile Range). Dots are outlier cases. Bar plots show change from pre to post measurements for the response variable.

Fig. 3 Change in BAI according to diagnosis



The box designates the median and the upper and lower quartiles. Whiskers/lines represent the max. and min. observations within the range of the 1. and 3. quartiles +/- 1.5\*IQR (InterQuartile Range). Dots are outlier cases. Bar plots show change from pre to post measurements for the response variable.

Fig. 4 Change in SES according to diagnosis